

Observations of Pearson-Readhead Survey  
Sources using VSOP and the EVN

Tingay, S.J., Preston, R.A., Piner, B.G.,  
Lister, M.L., Pearson, T.J., Readhead, A.C.S.,  
Murphy, D.W., Hirabayashi, H., Kobayashi,  
H., and Inoue, M.

We present high resolution images, preliminary analysis, and interpretation from VSOP space VLBI observations of Pearson-Readhead survey sources. We will highlight several of the sources that have been observed using a combination of the HALCA spacecraft and the EVN.

*astrophysics*

4<sup>th</sup> EVN/JIVE Symposium - October 1998


## INTRODUCTION

- The VSOP mission uses an Earth-orbiting antenna (HALCA; launched 1997 February) and ground-based facilities (radio telescopes, tracking stations, and correlators) to form a VLBI array with maximum baselines of approximately 30,000 km and  $u$ - $v$  coverage suitable for imaging.
- Observations at 5 GHz yield maximum resolution of approximately 0.2 mas, a good match to 15 GHz observations on the ground.
- For the 1<sup>st</sup> VSOP Announcement of Opportunity we proposed observations of sources from the complete Pearson-Readhead (PR) sample.
- The PR sample consists of 65 extragalactic radio sources with:
  - $\delta > 35^\circ$
  - $|b| > 10^\circ$
  - $S_{total}(5GHz) > 1.3$  Jy
- To select our sample of 31 sources from the PR sources we add:
  - $S_{>6000km}(5GHz) > 0.4$  Jy
- To date we have completed observations of 18/31 sources. Each observation consists of data from one HALCA orbit (6 hrs) and an array of ground radio telescopes. **For 6 of these observations EVN antennas have been used.**

## RESULTS

- Sources observed with VSOP/EVN observations include:
  - 1954+513
  - 2021+614
  - MKN501
  - 1642+690
  - 3C345
- We aim to measure accurate brightness temperatures ( $B_T$ ) for the PR sample.
  - $B_T$  sensitivity  $\propto D^2$
  - Ground-based observations are only sensitive to  $B_T \sim 10^{12}$  K  $\equiv$  nominal inverse Compton limit
  - Space VLBI observations are **required** to detect  $B_T > 10^{12}$  K
  - $B_T \propto \delta$
- What is the brightness temperature distribution for a complete sample of radio sources? Is there a characteristic maximum brightness temperature? How does it compare to the nominal  $10^{12}$  K inverse Compton limit.

Clean map. Array: EJMNSTW EJMNSTW EGMNSTW EGMNSW EGMNRSW EMNRSW  
1642+690 at 4.970 GHz 1998 May 31




Map center: RA: 16 42 07.849, Dec: +68 56 39.756 (2000.0)

Map peak: 0.794 Jy/beam

Contours %: -0.5 0.5 1 2 4 8 16 32 64


Beam FWHM: 6.19 x 3.99 (mas) at 74.3°

Clean map. Array: EJMNSTW EJMNSTW EGMNSTW EGMNSW EGMNRSW EMNRS  
1642+690 at 4.970 GHz 1998 May 31




Map center: RA: 16 42 07.849, Dec: +68 56 39.756 (2000.0)  
Map peak: 0.435 Jy/beam  
Contours %: -2 2 4 8 16 32 64  
Beam FWHM: 0.507 x 0.229 (mas) at  $-10.5^\circ$

1642+690 at 4.970 GHz 1998 May 31




Clean map. Array: EGMNORTWN  
1954+513 at 4.962 GHz 1997 Nov 10



Map center: RA: 19 55 42.738, Dec: +51 31 48.546 (2000.0)  
Map peak: 0.807 Jy/beam  
Contours %: -1 1 2 4 8 16 32 64  
Beam FWHM: 8.62 x 3.06 (mas) at 54.9°

Clean map. Array: EGMNORTWN  
1954+513 at 4.970 GHz 1997 Nov 10




Right Ascension (mas)


Map center: RA: 19 55 42.738, Dec: +51 31 48.546 (2000.0)  
Map peak: 0.313 Jy/beam  
Contours %: -5 5 10 20 40 80  
Beam FWHM: 0.424 x 0.254 (mas) at  $-40.2^\circ$



1954+513 at 4.962 GHz 1997 Nov 10




Clean map. Array: BEFHKLMMNNNOOPST  
2021+614 at 4.970 GHz 1997 Nov 07




Map center: RA: 20 22 06.682, Dec: +61 36 58.805 (2000.0)  
Map peak: 1.18 Jy/beam  
Contours %: -0.5 0.5 1 2 4 8 16 32 64  
Beam FWHM: 1.99 x 0.924 (mas) at -50.1°

Clean map. Array: BEFHKLMMNNNOOPSTYT  
2021+614 at 4.962 GHz 1997 Nov 07




Map center: RA: 20 22 06.682, Dec: +61 36 58.807 (2000.0)  
Map peak: 0.367 Jy/beam  
Contours %: -1 1 2 4 8 16 32 64  
Beam FWHM: 0.447 x 0.234 (mas) at -50.4°

2021+614 at 4.962 GHz 1997 Nov 07



Pearson-Readhead Source Brightness Temperatures (Gaussian)



Pearson-Readhead Sources Brightness Temperatures (Optically Thick Disk)

